MICHIGAN DEPARTMENT OF NATURAL RESOURCES

INTEROFFICE COMMUNICATION

January 25, 1993

TO: Petroleum hydrocarbon files

FROM: Gary Butterfield

SUBJECT: CAS numbers which can use the ITSL based on the NIOSH refined

petroleum products REL

A search for review documents on petroleum hydrocarbons located the 1977 NIOSH document for an REL of 350 mg/m3 for refined petroleum solvents. This document establishes an REL for various refined fractions and describes those fractions (petroleum ether, rubber solvent, VM & P naphtha, mineral spirits, and stoddard solvent) as having hydrocarbon chain lengths C5 to C12, a boiling point between 30 and 210 degrees Celsius, and being composed of less than 20 percent aromatic hydrocarbons. The chemical names and descriptions from the EPA ToSCA inventory and CAS-on-line were consulted to determine which CAS numbers might be covered by these three criteria, and could therefore, have an ITSL of 3500 ug/m3 (eight hour averaging time) established, as based on one percent of the NIOSH REL.

CAS No.	ToSCA Chemical name
8030-30-6	naphtha
8032-32-4	ligroine
8052-41-3	stoddard solvent
64742-48-9	naphtha (petroleum) hydrotreated heavy
64742-88-7	solvent naphtha (petroleum) medium aliphatic
64742-89-8	solvent naphtha (petroleum) light aliphatic

In addition to the above fractions, CAS # 64741-41-9 and ToSCA chemical name: naphtha (petroleum) heavy straight-run may also have the ITSL of 3500 ug/m3, with eight hour averaging, applied to it if it can be verified that the fraction contains less than 20 % aromatics. If the above materials are present in any combination, the ITSL applies to the total concentration of all those materials present - combined. As these petroleum hydrocarbon materials can be expected to impact the same organ systems because of their similarity, an additive effect is expected. Attached are printouts from CAS-on-line which contain CAS name, definition, and synonyms for each of the above petroleum fractions.

References:

- ACGIH. 1992. Threshold limit values for chemical substances and physical agents and biological exposure indices.
- EPA. 1979. ToSCA Inventory Appendix A.
- NIOSH. 1977. Criteria for a recommended standard ... occupational exposure to refined petroleum solvents. NIOSH 77-192.